in DC., Prodr. **11**: 92 (1847)—Wight, Icon. Pl. Ind. Or. **4**: t. 1490 (1850)—T. Anders. in Journ. Linn. Soc. **9**: 456 (1867), p.p.—C.B. Clarke in Fl. Brit. Ind. **4**: 407 (1884)—Matsum. et Hayata, Enum. Pl. Formos. 290 (1906)—Yamazaki in Fl. E. Himal. 302 (1966)—Fl. Hainan. **3**: 545, f. 919 (1974)—Icon. Cormophyt. Sin. **4**: 154, f. 5722 (1975).

Ruellia ringens L., Sp. Pl. **2**: 635 (1753), quoad specim. in Herb. Linn. tantum.

Ruellia salicifolia Vahl, Symb. **3**: 84 (1794)—Roxb., Fl. Ind. ed. 2, **3**: 50 (1832).

Justicia lancea Thunb. in Trans. Linn. Soc. **2**: 338 (1794).

Hygrophila angustifolia R. Brown, Prodr. Fl. Nov. Holl. 479 (1810).

Hygrophila lancea (Thunb.) Miquel in Ann. Mus. Bot. Lugd.-Bat. **2**: 123 (1865)—Matsum., Ind. Pl. Jap. **2**(2): 580 (1912)—Hara, Enum. Spermat. Jap. **1**: 294 (1949)—Kitamura et al., Col. Ill. Herb. Pl. Jap. **1**: 118 (1957)—Ohwi, Fl. Jap. ed. rev. 1229 (1965)—Walker, Fl. Okinawa 952 (1976).

Hygrophila pogonacalyx Hayata, Icon. Pl. Formos. **9**: 81 (1920).

Distr. India, Himalaya, Indo-China, east to S. China, Taiwan and S. & C. Japan, and Malaysia.

Very variable in the shape and size of leaves and the hairiness of leaves, bracts, calyces and corollae. In Japan it occurs only a form with almost glabrous lanceolate leaves, and it has always been treated as *H. lancea*. But the same form is also found in Himalaya and other regions. Both specific epithets, 'salicifolia' and 'lancea', were published in 1794, and I adopt 'salicifolia' which is well-known in the wider districts.

□木村陽二郎氏 Dr. Yojiro KIMURA

本誌編集員である木村陽二郎博士はフランス政府からパルムザカデミック勲章 (Palmes Académiques) のシュヴァアリエ章 (Chevalier) を1977年8月22日付で授与され，その伝達式が本年7月7日東京のフランス大使館で行われました。この勲章は1808年ナポレオンがフランス帝国大学を創立したのと同時に制定された由緒あるもので，学識ある教育の功労者に与えられ現在に至っています。木村博士は1954年ちょうど国際植物学会議がパリーで開かれた時に国立自然史博物館で一年間植物分類学および生物学史を研究され，1957年12月19日には同館の通信会員に選ばれました。また日仏会館会員，日仏生物学会幹事として日仏文化の交流発展に尽され，その功績が認められて今回この勲章を受けた最初の日本植物学者となられたことは誠に喜ばしいことで，ここに心からお祝を申し上げます。 (原 寛 Hiroshi HARA)